

# JINAZO®



## Product Introduction:

Active Ingredient	Content & Formulation
Azoxystrobin 3% + Jingtangmycin A 9%	12%WP

## Product Introduction:

Jingtangmycin A is an antibiotic fungicide that can interfere with the normal growth of pathogenic bacteria cells. Azoxystrobin is a mitochondrial respiration inhibitor that can block mitochondrial respiration, making it unable to produce ATP, and cause cell death. It has protection, treatment, eradication, infiltration and systemic activity, and inhibits spore germination and hypha growth. This formulation has good control effects on rice sheath blight.




## Advantage:


1. This product is an antibiotic compound formulation with a unique formula, high safety and low toxicity.
2. This product has the dual effect of prevention and treatment, and has good control effects on rice sheath blight and false smut. It can be effective for a long duration.

## Applicable Crops:


Rice, Grapes




Rice




Rice



Rice




Grapes




Grapes

## Targets:

Rice sheath blight, and false smut



Rice sheath blight



Rice false smut



Grape downy mildew



Rice false smut



Grape downy mildew

## Uses and Recommendations:

Crops	Targets	Dosage	Application method
Rice	Sheath blight	750-1050 g/ha	Spray
Rice	False smut	1500-2250 g/ha	Spray
Rice	Rice blast	1500-2250 g/ha	Spray
Grape	White rot	1000-1500 times diluted with water	Spray
Grape	Anthracoze	1000-1500 times diluted with water	Spray
Grape	Downy mildew	1000-1500 times diluted with water	Spray

1. The pesticide should be applied once during the rice breakage period and the full heading period, and sprayed evenly with water according to the recommended dose; the spraying should be comprehensive to prevent spray leakage, and the water depth of the paddy field should be maintained at 3-6 cm when spraying.
2. After application of this product on the rice, it should be at least 30 days apart from the application to making the harvest. This product is limited to two applications per crop season.
3. Do not apply in windy days or when rainfall is expected within one hour.

## Cautions:

1. Do not mix with alkaline pesticides and other substances.
2. Wear protective clothing, protective gloves, masks, etc. when using this product to avoid skin contact and mouth and nose inhalation.